Methods

Study Selection

We sought papers published by investigators referencing datasets from The Genetic Association Information Network (GAIN). We searched through papers included in The GAIN Data Access Committee (GAIN DAC) annual reports of investigators requesting access to GAIN genotyping data.

Only publications written by study investigators acknowledging use of GAIN datasets were included in the Publications Using GAIN Genotyping Data bibliography. These publications were scanned for search terms relating to GAIN or dbGaP (ie: *GAIN, dbGaP, data accession number, etc*).

Publications Using GAIN Genotyping Data

1. Jian XQ, Wang KS, Wu TJ, Hillhouse JJ, Mullersman JE. Association of ADAM10 and CAMK2A Polymorphisms with Conduct Disorder: Evidence from Family-Based Studies. J Abnorm Child Psychol 39: 773-82. 2011.

2. Allen AS, Satten GA. Control for confounding in case-control studies using the stratification score, a retrospective balancing score. Am J Epidemiol 173: 752-60. 2011.

3. Cichon S, Muhleisen TW, Degenhardt FA, Mattheisen M, Miro X, Strohmaier J, Steffens M, Meesters C, Herms S, Weingarten M, Priebe L, Haenisch B, Alexander M, Vollmer J, Breuer R, Schmal C, et al. Genome-wide association study identifies genetic variation in neurocan as a susceptibility factor for bipolar disorder. Am J Hum Genet 88: 372-81. 2011.

4. Zhang F, Chen Q, Ye T, Lipska BK, Straub RE, Vakkalanka R, Rujescu D, St Clair D, Hyde TM, Bigelow L, Kleinman JE, Weinberger DR. Evidence of Sex-Modulated Association of ZNF804A with Schizophrenia. Biol Psychiatry 69: 914-7. 2011.

5. Kendler KS, Kalsi G, Holmans PA, Sanders AR, Aggen SH, Dick DM, Aliev F, Shi J, Levinson DF, Gejman PV. Genomewide Association Analysis of Symptoms of Alcohol Dependence in the Molecular Genetics of Schizophrenia (MGS2) Control Sample. Alcohol Clin Exp Res 35: 963-975. 2011.

6. Belmonte Mahon P, Pirooznia M, Goes FS, Seifuddin F, Steele J, Lee PH, Huang J, Hamshere ML, Depaulo JR, Jr., Kelsoe JR, Rietschel M, Nothen M, Cichon S, Gurling H, Purcell S, Smoller JW, Craddock N, Schulze TG, McMahon FJ, Potash JB, Zandi PP. Genome-wide association analysis of age at onset and psychotic symptoms in bipolar disorder. Am J Med Genet B Neuropsychiatr Genet 156: 370-378. 2011.

Last updated: July 22nd, 2011

7. Levinson DF, Duan J, Oh S, Wang K, Sanders AR, Shi J, Zhang N, Mowry BJ, Olincy A, Amin F, Cloninger CR, Silverman JM, Buccola NG, Byerley WF, Black DW, Kendler KS, Freedman R, Dudbridge F, Pe'er I, Hakonarson H, Bergen SE, Fanous AH, Holmans PA, Gejman PV. Copy number variants in schizophrenia: confirmation of five previous findings and new evidence for 3q29 microdeletions and VIPR2 duplications. Am J Psychiatry 168: 302-16. 2011.

8. Sun J, Wan C, Jia P, Fanous AH, Kendler KS, Riley BP, Zhao Z. Application of systems biology approach identifies and validates GRB2 as a risk gene for schizophrenia in the Irish Case Control Study of Schizophrenia (ICCSS) sample. Schizophr Res 125: 201-8. 2011.

9. Hartz SM, Lin P, Edenberg HJ, Xuei X, Rochberg N, Saccone S, et al. Genetic association of bipolar disorder with the beta3 nicotinic receptor subunit gene. Psychiatr Genet. 2010. Available online only.

10. Machado-Vieira R, Pivovarova NB, Stanika RI, Yuan P, Wang Y, Zhou R, Zarate CA, Jr., Drevets WC, Brantner CA, Baum A, Laje G, McMahon FJ, Chen G, Du J, Manji HK, Andrews SB. The Bcl-2 gene polymorphism rs956572AA increases inositol 1,4,5-trisphosphate receptor-mediated endoplasmic reticulum calcium release in subjects with bipolar disorder. Biol Psychiatry 69: 344-52. 2010.

11. Kornum BR, Kawashima M, Faraco J, Lin L, Rico TJ, Hesselson S, Axtell RC, Kuipers H, Weiner K, Hamacher A, et al. Common variants in P2RY11 are associated with narcolepsy. Nat Genet 43: 66-71. 2010.

12. Peterson RE, Maes HH, Holmans P, Sanders AR, Levinson DF, Shi J, Kendler KS, Gejman PV, Webb BT. Genetic risk sum score comprised of common polygenic variation is associated with body mass index. Hum Genet 129: 221-30. 2010.

13. Glessner JT, Wang K, Sleiman PM, Zhang H, Kim CE, Flory JH, Bradfield JP, Imielinski M, Frackelton EC, Qiu H, Mentch F, Grant SF, Hakonarson H. Duplication of the SLIT3 locus on 5q35.1 predisposes to major depressive disorder. PLoS One. 2010. Available online only.

14. Valeyev NV, Hundhausen C, Umezawa Y, Kotov NV, Williams G, Clop A, Ainali C, Ouzounis C, Tsoka S, Nestle FO. A systems model for immune cell interactions unravels the mechanism of inflammation in human skin. PLoS Comput Biol 6. 2010. Available online only.

15. Padmanabhan S, Melander O, Johnson T, Di Blasio AM, Lee WK, Gentilini D, et al. Genome-wide association study of blood pressure extremes identifies variant near UMOD associated with hypertension. PLoS Genet. 2010; Available online only.

Last updated: July 22nd, 2011

16. Wray NR, Pergadia ML, Blackwood DH, Penninx BW, Gordon SD, Nyholt DR, et al. Genome-wide association study of major depressive disorder: new results, meta-analysis, and lessons learned. Mol Psychiatry. 2010. Available online only.

17. Zawistowski M, Gopalakrishnan S, Ding J, Li Y, Grimm S, Zollner S. Extending rare-variant testing strategies: analysis of noncoding sequence and imputed genotypes. Am J Hum Genet 87: 604-17. 2010.

18. Pan Y, Wang KS, Aragam N. NTM and NR3C2 polymorphisms influencing intelligence: family-based association studies. Prog Neuropsychopharmacol Biol Psychiatry. 2011; 35: 154-60.

19. Huang N, Lee I, Marcotte EM, Hurles ME. Characterising and predicting haploinsufficiency in the human genome. PLoS Genet. 2010 Available online only.

20. Heid IM, Jackson AU, Randall JC, Winkler TW, Qi L, Steinthorsdottir V, Thorleifsson G, Zillikens MC, Speliotes EK, et al. Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. Nat Genet 42: 949-60. 2010.
21. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. Nat Genet. 2010; 42: 937-48.

22. Klok MD, Vreeburg SA, Penninx BW, Zitman FG, de Kloet ER, Derijk RH. Common functional mineralocorticoid receptor polymorphisms modulate the cortisol awakening response: Interaction with SSRIs. Psychoneuroendocrinology. 2010; 4: 484-94.

23. Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F, Willer CJ, Jackson AU, Vedantam S, Raychaudhuri S, , et al. Hundreds of variants clustered in genomic loci and biological pathways affect human height. Nature 467: 832-8. 2010.

24. Knight J, Rochberg NS, Saccone SF, Nurnberger JI, Jr., Rice JP. An investigation of candidate regions for association with bipolar disorder. Am J Med Genet B Neuropsychiatr Genet 153B: 1292-7. 2010.

25. Soranzo N, Sanna S, Wheeler E, Gieger C, Radke D, Dupuis J, et al. Common variants at 10 genomic loci influence hemoglobin A(C) levels via glycemic and nonglycemic pathways. Diabetes. 2010; 59: 3229-39.

26. Chen X, Lee G, Maher BS, Fanous AH, Chen J, Zhao Z, et al. GWA study data mining and independent replication identify cardiomyopathy-associated 5

(CMYA5) as a risk gene for schizophrenia. Mol Psychiatry. 2010. Available online only.

27. Carvalho BS, Louis TA, Irizarry RA. Quantifying uncertainty in genotype calls. Bioinformatics. 26: 242-9. 2010.

28. Bray SM, Mulle JG, Dodd AF, Pulver AE, Wooding S, Warren ST. Signatures of founder effects, admixture, and selection in the Ashkenazi Jewish population. Proc Natl Acad Sci U S A. 2010; 107: 16222-7.

29. Neale BM, Medland SE, Ripke S, Asherson P, Franke B, Lesch KP, et al. Meta-analysis of genome-wide association studies of attention-deficit/hyperactivity disorder. J Am Acad Child Adolesc Psychiatry. 2010; 49: 884-97.

30. Dao DT, Mahon PB, Cai X, Kovacsics CE, Blackwell RA, Arad M, Shi J, Zandi PP, O'Donnell P, Knowles JA, Weissman MM, Coryell W, Scheftner WA, Lawson WB, Levinson DF, Thompson SM, Potash JB, Gould TD. Mood disorder susceptibility gene CACNA1C modifies mood-related behaviors in mice and interacts with sex to influence behavior in mice and diagnosis in humans. Biol Psychiatry 68: 801-10. 2010.

31. Mick E, Todorov A, Smalley S, Hu X, Loo S, Todd RD, Biederman J, Byrne D, Dechairo B, Guiney A, McCracken J, McGough J, Nelson SF, Reiersen AM, Wilens TE, Wozniak J, Neale BM, Faraone SV. Family-based genome-wide association scan of attention-deficit/hyperactivity disorder. J Am Acad Child Adolesc Psychiatry 49: 898-905. 2010.

32. Mulle JG, Dodd AF, McGrath JA, Wolyniec PS, Mitchell AA, Shetty AC, et al. Microdeletions of 3q29 confer high risk for schizophrenia. Am J Hum Genet. 2010; 87: 229-36.

33. Agrawal A, Nurnberger JI, Jr., Lynskey MT.Cannabis involvement in individuals with bipolar disorder. Psychiatry Res 185: 459-61. 2010.

34. Jia P, Wang L, Meltzer HY, Zhao Z. Common variants conferring risk of schizophrenia: a pathway analysis of GWAS data. Schizophr Res. 2010; 122: 38-42.

35. Eijgelsheim M, Newton-Cheh C, Sotoodehnia N, de Bakker PI, Muller M, Morrison AC, et al. Genome-wide association analysis identifies multiple loci related to resting heart rate. Hum Mol Genet. 2010; 19: 3885-94.

36. Scharpf RB, Ruczinski I, Carvalho B, Doan B, Chakravarti A, Irizarry RA. A multilevel model to address batch effects in copy number estimation using SNP arrays. *Biostatistics*. 2011; 12: 33-50.

37. Sun J, Jia P, Fanous AH, van den Oord E, Chen X, Riley BP, et al. Schizophrenia gene networks and pathways and their applications for novel candidate gene selection. PLoS One. 2010 Available online only.

38. Pluzhnikov A, Below JE, Konkashbaev A, Tikhomirov A, Kistner-Griffin E, Roe CA, et al. Spoiling the whole bunch: quality control aimed at preserving the integrity of high-throughput genotyping. Am J Hum Genet. 2010; 87: 123-8.

39. Demirkan A, Penninx BW, Hek K, Wray NR, Amin N, Aulchenko YS, et al. Genetic risk profiles for depression and anxiety in adult and elderly cohorts. Mol Psychiatry. 2010. Available online only.

40. Alvarado DM, Aferol H, McCall K, Huang JB, Techy M, Buchan J et al. Familial isolated clubfoot is associated with recurrent chromosome 17q23.1q23.2 microduplications containing TBX4. Am J Hum Genet; 87: 154-160. 2010.

41. Oedegaard KJ, Greenwood TA, Johansson S, Jacobsen KK, Halmoy A, Fasmer OB, Akiskal HS, Haavik J, Kelsoe JR.A genome-wide association study of bipolar disorder and comorbid migraine. Genes Brain Behav 9: 673-80. 2010.

42. Sanders AR, Levinson DF, Duan J, Dennis JM, Li R, Kendler KS, Rice JP, Shi J, Mowry BJ, Amin F, Silverman JM, Buccola NG, Byerley WF, Black DW, Freedman R, Cloninger CR, Gejman PV. The Internet-based MGS2 control sample: self report of mental illness. Am J Psychiatry 167: 854-65. 2010.

43. Jackson KJ, Chen Q, Chen J, Aggen SH, Kendler KS, Chen X. Association of the histidine-triad nucleotide-binding protein-1 (HINT1) gene variants with nicotine dependence. Pharmacogenomics J. 2010. Available online only.

44. Pongpanich M, Sullivan PF, Tzeng JY. A quality control algorithm for filtering SNPs in genome-wide association studies. Bioinformatics 26: 1731-7. 2010.

45. Glessner JT, Reilly MP, Kim CE, Takahashi N, Albano A, Hou C, et al. Strong synaptic transmission impact by copy number variations in schizophrenia. Proc Natl Acad Sci U S A. 2010; 107: 10584-9.

46. Thorgeirsson TE, Gudbjartsson DF, Surakka I, Vink JM, Amin N, Geller F, et al. Sequence variants at CHRNB3-CHRNA6 and CYP2A6 affect smoking behavior. Nat Genet. 2010; 42: 448-53.

47. Tobacco and Genetics Consortium. Genome-wide meta-analyses identify multiple loci associated with smoking behavior. Nat Genet. 2010; 42: 441-7.

48. Zou F, Lee S, Knowles MR, Wright FA. Quantification of population structure using correlated SNPs by shrinkage principal components. Hum Hered. 2010; 70: 9-22.

49. Chambers JC, Zhang W, Lord GM, van der Harst P, et al. Genetic loci influencing kidney function and chronic kidney disease. Nat Genet 42: 373-5. 2010.

50. Freathy RM, Mook-Kanamori DO, Sovio U, Prokopenko I, Timpson NJ, Berry DJ, et al. Variants in ADCY5 and near CCNL1 are associated with fetal growth and birth weight. Nat Genet. 2010; 42: 430-5.

51. Bosker FJ, Hartman CA, Nolte IM, Prins BP, Terpstra P, Posthuma D, van Veen T, Willemsen G, Derijk RH, de Geus EJ, Hoogendijk WJ, Sullivan PF, Penninx BW, Boomsma DI, Snieder H, Nolen WA. Poor replication of candidate genes for major depressive disorder using genome-wide association data. Mol Psychiatry. 2010; 5: 516-32.

52. Cannon DM, Klaver JK, Gandhi SK, Solorio G, Peck SA, Erickson K, JS NA, Eckelman WC, Furey ML, Sahakian BJ, McMahon FJ, Drevets WC. Genetic variation in cholinergic muscarinic-2 receptor gene modulates M2 receptor binding in vivo and accounts for reduced binding in bipolar disorder. Mol Psychiatry 16: 407-18. 2010.

53. Liu Y, Blackwood DH, Caesar S, de Geus EJ, Farmer A, Ferreira MA, Ferrier IN, Fraser C, Gordon-Smith K, Green EK, Grozeva D, Gurling HM, Hamshere ML, Heutink P, et al. Meta-analysis of genome-wide association data of bipolar disorder and major depressive disorder. Mol Psychiatry 16: 2-4. 2010.

54. Lasky-Su J, Won S, Mick E, Anney RJ, Franke B, Neale B, *et al.* On genome-wide association studies for family-based designs: an integrative analysis approach combining ascertained family samples with unselected controls. Am J Hum Genet. 2010; 86: 573-80.

55. Gejman PV, Sanders AR, Duan J. The role of genetics in the etiology of schizophrenia. Psychiatr Clin North Am 33: 35-66. 2010.

56. O'Dushlaine C, Kenny E, Heron E, Donohoe G, Gill M, Morris D, *et al.* Molecular pathways involved in neuronal cell adhesion and membrane scaffolding contribute to schizophrenia and bipolar disorder susceptibility. Mol Psychiatry. 2010; 3: 286-92.

57. Shi J, Potash JB, Knowles JA, Weissman MM, Coryell W, Scheftner WA, Lawson WB, Depaulo JR, Jr., Gejman PV, Sanders AR, Johnson JK, Adams P, Chaudhury S, Jancic D, Evgrafov O, Zvinyatskovskiy A, Ertman N, Gladis M, Neimanas K, Goodell M, Hale N, Ney N, Verma R, Mirel D, Holmans P, Levinson DF. Genome-wide association study of recurrent early-onset major depressive disorder. Mol Psychiatry 2010; 16: 193-201.

58. McMahon FJ, Akula N, Schulze TG, Muglia P, Tozzi F, Detera-Wadleigh SD, *et al.* Meta-analysis of genome-wide association data identifies a risk locus for major mood disorders on 3p21.1. Nat Genet. 2010; 42: 128-31.

59. Ruano D, Abecasis GR, Glaser B, Lips ES, Cornelisse LN, de Jong AP, *et al.* Functional gene group analysis reveals a role of synaptic heterotrimeric G proteins in cognitive ability. Am J Hum Genet. 2010; 86: 113-25.

60. Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, Wheeler E, et al. New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. Nat Genet 42: 105-16. 2010.

61. Shyn SI, Shi J, Kraft JB, Potash JB, Knowles JA, Weissman MM, Garriock HA, Yokoyama JS, McGrath PJ, Peters EJ, Scheftner WA, Coryell W, Lawson WB, Jancic D, Gejman PV, Sanders AR, Holmans P, Slager SL, Levinson DF, Hamilton SP. Novel loci for major depression identified by genome-wide association study of sequenced treatment alternatives to relieve depression and meta-analysis of three studies. Mol Psychiatry. 2009; 2: 202-15. .

62. Fullerton JM, Donald JA, Mitchell PB, Schofield PR. Two-dimensional genome scan identifies multiple genetic interactions in bipolar affective disorder. Biol Psychiatry. 2009. 67: 478-86.

63. Bochukova EG, Huang N, Keogh J, Henning E, Purmann C, Blaszczyk K, Saeed S, Hamilton-Shield J, Clayton-Smith J, O'Rahilly S, Hurles ME, Farooqi IS. Large, rare chromosomal deletions associated with severe early-onset obesity, Nature 2009; 463: 666-70.

64. Wallace C, Smyth DJ, Maisuria-Armer M, Walker NM, Todd JA, Clayton DG. The imprinted dlk1-meg3 gene region on chromosome 14q32.2 alters susceptibility to type 1 diabetes. Nat Genet 2009; 42:68-71.

65. Tsaih SW, Pezzolesi MG, Yuan R, Warram JH, Krolewski AS, Korstanje R. Genetic analysis of albuminuria in aging mice and concordance with loci for human diabetic nephropathy found in a genome-wide association scan, Kidney Int 2009; 77:201-10.

66. Paterson AD, Waggott D, Boright AP, Hosseini SM, Shen E, Sylvestre MP, Wong I, Bharaj B, Cleary PA, Lachin JM, Below JE, Nicolae D, Cox NJ, Canty AJ, Sun L, Bull SB. A genome-wide association study identifies a novel major locus for glycemic control in type 1 diabetes, as measured by both hba1c and glucose, Diabetes 2009; 59: 332-4.

67. McCarthy SE, Makarov V, Kirov G, Addington AM, McClellan J, Yoon S, Perkins DO, Dickel DE, Kusenda M, Krastoshevsky O, Krause V, Kumar RA, Grozeva D, Malhotra D, Walsh T, Zackai EH, Kaplan P, Ganesh J, Krantz ID,

Spinner NB, Roccanova P, Bhandari A, Pavon K, Lakshmi B, Leotta A, Kendall J, Lee YH, Vacic V, Gary S, Iakoucheva LM, Crow TJ, Christian SL, Lieberman JA, Stroup TS, Lehtimaki T, Puura K, Haldeman-Englert C, Pearl J, Goodell M, Willour VL, Derosse P, Steele J, Kassem L, Wolff J, Chitkara N, McMahon FJ, Malhotra AK, Potash JB, Schulze TG, Nothen MM, Cichon S, Rietschel M, Leibenluft E, Kustanovich V, Lajonchere CM, Sutcliffe JS, Skuse D, Gill M, Gallagher L, Mendell NR, Craddock N, Owen MJ, O'Donovan MC, Shaikh TH, Susser E, Delisi LE, Sullivan PF, Deutsch CK, Rapoport J, Levy DL, King MC, Sebat J. Microduplications of 16p11.2 are associated with schizophrenia, Nat Genet 2009; 41:1223-1227.

68. Paterson AD, Lopes-Virella MF, Waggott D, Boright AP, Hosseini SM, Carter RE, Shen E, Mirea L, Bharaj B, Sun L, Bull SB. Genome-wide association identifies the abo blood group as a major locus associated with serum levels of soluble e-selectin. Arterioscler Thromb Vasc Biol 2009; 29:1958-1967.

69. Trevino LR, Yang W, French D, Hunger SP, Carroll WL, Devidas M, Willman C, Neale G, Downing J, Raimondi SC, Pui CH, Evans WE, Relling MV. Germline genomic variants associated with childhood acute lymphoblastic leukemia, Nat Genet 2009; 41: 1001-1005.

70. Yoo YJ, Bull SB, Paterson AD, Waggott D, Sun L. Were genome-wide linkage studies a waste of time? Exploiting candidate regions within genome-wide association studies, Genet Epidemiol 2009; 34:107-118.

71. Sun J, Jia P, Fanous AH, Webb BT, van den Oord EJ, Chen X, Bukszar J, Kendler KS, Zhao Z. A multi-dimensional evidence-based candidate gene prioritization approach for complex diseases-schizophrenia as a case, Bioinformatics Jul 16 2009; 25: 2595-6602.

72. Quaranta, M, Burden, AD, Griffiths, CE, Worthington, J, Barker, JN, Trembath, RC, Capon, F. Differential contribution of CDKAL1 variants to psoriasis, Crohn's disease and type II diabetes, Genes Immun 2009; 10: 654-8.

73. Shi, J, Levinson, DF, Duan, J, Sanders, AR, Zheng, Y, Pe'er, I, Dudbridge, F, Holmans, PA, Whittemore, AS, Mowry, BJ, Olincy, A, Amin, F, Cloninger, CR, Silverman, JM, Buccola, NG, Byerley, WF, Black, DW, Crowe, RR, Oksenberg, JR, Mirel, DB, Kendler, KS, Freedman, R, Gejman, PV. Common variants on chromosome 6p22.1 are associated with schizophrenia, Nature 2009; 460: 753-757.

74. Estrada, K, Krawczak, M, Schreiber, S, van Duijn, K, Stolk, L, van Meurs, JB, Liu, F, Penninx, BW, Smit, JH, Vogelzangs, N, Hottenga, JJ, Willemsen, G, de Geus, EJ, Lorentzon, M, von Eller-Eberstein, H, Lips, P, Schoor, N, Pop, V, de Keijzer, J, Hofman, A, Aulchenko, YS, Oostra, BA, Ohlsson, C, Boomsma, DI, Uitterlinden, AG, van Duijn, CM, Rivadeneira, F, Kayser, M. A genome-wide

association study of northwestern Europeans involves the C-type natriuretic peptide signaling pathway in the etiology of human height variation, Hum Mol Genet 2009; 18: 3516-3524.

75. Hao, K, Chudin, E, McElwee, J, Schadt, EE. Accuracy of genome-wide imputation of untyped markers and impacts on statistical power for association studies, BMC Genet 2009; 10: 27.

76. Barrett, JC, Clayton, DG, Concannon, P, Akolkar, B, Cooper, JD, Erlich, HA, Julier, C, Morahan, G, Nerup, J, Nierras, C, Plagnol, V, Pociot, F, Schuilenburg, H, Smyth, DJ, Stevens, H, Todd, JA, Walker, NM, Rich, SS. Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes, Nat Genet 2009; 41: 703-7.

77. Hallmayer, J, Faraco, J, Lin, L, Hesselson, S, Winkelmann, J, Kawashima, M, Mayer, G, Plazzi, G, Nevsimalova, S, Bourgin, P, Hong, SS, Honda, Y, Honda, M, Hogl, B, Longstreth, WT, Jr., Montplaisir, J, Kemlink, D, Einen, M, Chen, J, Musone, SL, Akana, M, Miyagawa, T, Duan, J, Desautels, A, Erhardt, C, Hesla, PE, Poli, F, Frauscher, B, Jeong, JH, Lee, SP, Ton, TG, Kvale, M, Kolesar, L, Dobrovolna, M, Nepom, GT, Salomon, D, Wichmann, HE, Rouleau, GA, Gieger, C, Levinson, DF, Gejman, PV, Meitinger, T, Young, T, Peppard, P, Tokunaga, K, Kwok, PY, Risch, N, Mignot, E. Narcolepsy is strongly associated with the T-cell receptor alpha locus, Nat Genet 2009; 41: 708-711.

78. Vink JM, Smit AB, de Geus EJ, Sullivan P, Willemsen G, Hottenga JJ, Smit JH, Hoogendijk WJ, Zitman FG, Peltonen L, Kaprio J, Pedersen NL, Magnusson PK, Spector TD, Kyvik KO, Morley KI, Heath AC, Martin NG, Westendorp RG, Slagboom PE, Tiemeier H, Hofman A, Uitterlinden AG, Aulchenko YS, Amin N, van Duijn C, Penninx BW, Boomsma DI. Genome-wide association study of smoking initiation and current smoking, Am J Hum Genet 2009; 84:367-379.

79. Schimmelmann, BG, Friedel, S, Nguyen, TT, Sauer, S, Ganz Vogel, CI, Konrad, K, Wilhelm, C, Sinzig, J, Renner, TJ, Romanos, M, Palmason, H, Dempfle, A, Walitza, S, Freitag, C, Meyer, J, Linder, M, Schafer, H, Warnke, A, Lesch, KP, Herpertz-Dahlman, B, Hinney, A, Hebebrand, J. Exploring the genetic link between RLS and ADHD, J Psychiatr Res 2009; 43: 941-945.

80. Pardo, L, Bochdanovits, Z, de Geus, E, Hottenga, JJ, Sullivan, P, Posthuma, D, Penninx, BW, Boomsma, D, Heutink, P. Global similarity with local differences in linkage disequilibrium between the Dutch and HapMap-CEU populations, Eur J Hum Genet 2009; 17: 802-810.

81. Aulchenko, YS, Ripatti, S, Lindqvist, I, Boomsma, D, Heid, IM, Pramstaller, PP, Penninx, BW, Janssens, AC, Wilson, JF, Spector, T, Martin, NG, Pedersen, NL, Kyvik, KO, Kaprio, J, Hofman, A, Freimer, NB, Jarvelin, MR, Gyllensten, U, Campbell, H, Rudan, I, Johansson, A, Marroni, F, Hayward, C, Vitart, V,

Jonasson, I, Pattaro, C, Wright, A, Hastie, N, Pichler, I, Hicks, AA, Falchi, M, Willemsen, G, Hottenga, JJ, de Geus, EJ, Montgomery, GW, Whitfield, J, Magnusson, P, Saharinen, J, Perola, M, Silander, K, Isaacs, A, Sijbrands, EJ, Uitterlinden, AG, Witteman, JC, Oostra, BA, Elliott, P, Ruokonen, A, Sabatti, C, Gieger, C, Meitinger, T, Kronenberg, F, Doring, A, Wichmann, HE, Smit, JH, McCarthy, MI, van Duijn, CM, Peltonen, L. Loci influencing lipid levels and coronary heart disease risk in 16 European population cohorts, Nat Genet 2008; 41: 47-55.

82. Prokopenko, I, Langenberg, C, Florez, JC, Saxena, R, Soranzo, N, Thorleifsson, G, Loos, RJ, Manning, AK, Jackson, AU, Aulchenko, Y, Potter, SC, Erdos, MR, Sanna, S, Hottenga, JJ, Wheeler, E, Kaakinen, M, Lyssenko, V, Chen, WM, Ahmadi, K, Beckmann, JS, Bergman, RN, Bochud, M, Bonnycastle, LL, Buchanan, TA, Cao, A, Cervino, A, Coin, L, Collins, FS, Crisponi, L, de Geus, E.J., Dehghan, A, Deloukas, P, Doney, AS, Elliott, P, Freimer, N, Gateva, V, Herder, C, Hofman, A, Hughes, TE, Hunt, S, Illig, T, Inouye, M, Isomaa, B, Johnson, T, Kong, A, Krestyaninova, M, Kuusisto, J, Laakso, M, Lim, N, Lindblad, U, Lindgren, CM, McCann, OT, Mohlke, KL, Morris, AD, Naitza, S, Orru, M, Palmer, CN, Pouta, A, Randall, J, Rathmann, W, Saramies, J, Scheet, P, Scott, LJ, Scuteri, A, Sharp, S, Sijbrands, E, Smit, JH, Song, K, Steinthorsdottir, V, Stringham, HM, Tuomi, T, Tuomilehto, J, Uitterlinden, AG, Voight, BF, Waterworth, D, Wichmann, HE, Willemsen, G, Witteman, JC, Yuan, X, Zhao, JH, Zeggini, E, Schlessinger, D, Sandhu, M, Boomsma, DI, Uda, M, Spector, TD, Penninx, BW, Altshuler, D, Vollenweider, P, Jarvelin, MR, Lakatta, E, Waeber, G, Fox, CS, Peltonen, L, Groop, LC, Mooser, V, Cupples, LA, Thorsteinsdottir, U, Boehnke, M, Barroso, I, Van Duijn, C, Dupuis, J, Watanabe, RM, Stefansson, K, McCarthy, MI, Wareham, NJ, Meigs, JB, Abecasis, GR. Variants in MTNR1B influence fasting glucose levels, Nat Genet 2008; 41: 77-81.

83. Jung, J, Sun, B, Kwon, D, Koller, DL, Foroud, TM. Allelic-based gene-gene interaction associated with quantitative traits, Genet Epidemiol 2009; 33: 332-343.

84. Gusev, A, Lowe, JK, Stoffel, M, Daly, MJ, Altshuler, D, Breslow, JL, Friedman, JM, Pe'er, I. Whole population, genome-wide mapping of hidden relatedness, Genome Res 2008; 19: 318-326.

85. Mulligan A, Richardson T, Anney RJ, Gill M. The Social Communication Questionnaire in a sample of the general population of school-going children. Ir J Med Sci 178: 193-9. 2008.

86. Payne, JL, MacKinnon, DF, Mondimore, FM, McInnis, MG, Schweizer, B, Zamoiski, RB, McMahon, FJ, Nurnberger, JI, Jr., Rice, JP, Scheftner, W, Coryell, W, Berrettini, WH, Kelsoe, JR, Byerley, W, Gershon, ES, DePaulo, JR, Jr., Potash, JB. Familial aggregation of postpartum mood symptoms in bipolar disorder pedigrees, Bipolar Disord 2008; 10: 38-44.

87. Goes, FS, Zandi, PP, Miao, K, McMahon, FJ, Steele, J, Willour, VL, Mackinnon, DF, Mondimore, FM, Schweizer, B, Nurnberger, JI, Jr., Rice, JP, Scheftner, W, Coryell, W, Berrettini, WH, Kelsoe, JR, Byerley, W, Murphy, DL, Gershon, ES, Bipolar Disorder Phenome, G, Depaulo, JR, Jr., McInnis, MG, Potash, JB. Mood-incongruent psychotic features in bipolar disorder: familial aggregation and suggestive linkage to 2p11-q14 and 13q21-33, Am J Psychiatry 2007; 164: 236-247.